46. A List of the Snakes of North-East Africa, from the Tropic to the Soudan and Somaliland, including Socotra. By G. A. BOULENGER, F.R.S., F.Z.S.*

[Received October 4, 1915; Read November 23, 1915.]

		х	

GEOGRAPHICAL :	Page
N.E. Africa, List of Snakes	641
SYSTEMATIC : Keys to the identification of the Snakes of N.E. Africa	641

With this list the Ophidian fauna of East and Central Africa is completed, only two divisions remaining to be dealt with, viz., West Africa north of the Congo, and North Africa.

Synopsis of the Families.

I. Worm-like	, with sma	ll inferior	mouth; e	eyes hidden	or visible	under th	e head-
shields, a	nd body co	vered with	uniform	imbricate :	scales abov	ve and ber	leath.

22 or more scales round middle of body; ocular not bordering the	
mouth; tail not or but little longer than broad	TYPHLOPIDÆ.
14 scales round middle of body	GLAUCONIID.E.
11. Mouth large, eyes exposed; body with enlarged shields be	eneath (except in
the marine genus <i>Hydrus</i>).	
Ventral shields much narrower than the body; supraocular, if	
distinct, broken up into two or more shields; scales on	
anterior part of body smooth	BOIDÆ.
Ventral shields at least nearly as broad as the body ; supraocular	
single; poison-fangs, if present, not in a very large sheath	Colubrid.e.
Ventral shields at least nearly as broad as the body, or upper sur-	
face of head covered with small scales; poisou-tangs in a very	
large sheath below the eye	VIPERID.E.

Family TYPHLOPIDÆ.

A single genus.

1. TYPHLOPS.

Schneid, Hist. Amph. ii. p. 339; Bouleng, Cat. Sn. i. p. 7.

Synopsis of the Species.

I. Snout rounded or with obtusely angular horizontal edge.

A. Rostral not more than one-third width of head; eyes distinct.

Snout rounded ; 24 scales round middle of body, the diameter of which is 37 to 50 times in total length	T. socotranus.
Snout much depressed; 22 scales round middle of body, the diameter of which is 20 to 30 times in total length	T. cuneirostris.
B. Rostral very broad.	
Rostral half width of head: 30 scales round middle of body, the diameter of which is 30 to 40 times in total length	T. blanfordii.
Rostral more than half width of head; 26 to 30 scales round	
middle of body, the diameter of which is 24 to 30 times in total length	T. punctatus.

* Published by permission of the Trustees of the British Museum.